

Comment on "Cerenkov radiation by neutrinos in a supernova core"

Subhendra Mohanty¹ and Sarira Sahu²

Theory Group, Physical Research Laboratory,
Navrangpura, Ahmedabad-380 009
India

It had been pointed out by Mohanty and Samal[1] that the helicity flipping Cerenkov process $\nu_L \rightarrow \nu_R + \gamma$ or $\gamma + \nu_L \rightarrow \nu_R$ could be an important cooling mechanism for the supernova core. Comparing the neutrino emissivity by the Cerenkov process with observations of SN1987A, a restrictive bound on the neutrino magnetic moment was established. Subsequently it was pointed out by Raffelt[2] that, this result was based on a numerical error in the calculation of the refractive index of the SN core and using the correct numbers it was shown that the photons in a SN core do not have a space-like dispersion relation, so the Cerenkov helicity flip process would not occur.

Here we show that the earlier estimate of refractive index was based on the thermodynamic formula for susceptibility which turns out to be invalid for real photons or plasmons even in the static limit. However an analysis of the dispersion relations of plasmons in an

ultrarelativistic plasma[3] shows that the longitudinal photon (plasmon) has a space-like branch and therefore in such a plasma the Cerenkov radiation of a plasmon is kinematically allowed[4]. We compute the neutrino emissivity of SN core by the Cerenkov helicity flip emission of a plasmon and subsequent escape of the ν_R . We show that the observations of neutrino flux from SN1987A put a constraint on the neutrino magnetic moment $\mu_\nu < 0.6 \times 10^{-11} \mu_B$.

The dispersion relation of a longitudinal photon (plasmon) is given by [3]

$$k^2 - \frac{3}{2}m_\gamma^2 \left[\frac{w}{k} \ln \left(\left| \frac{w+k}{w-k} \right| \right) - 2 \right] = 0. \quad (1)$$

Using this dispersion relation we find that in a supernova core there is a range of frequencies $(0.3 - 50) \text{ MeV}$ for which the refractive index $n = k/w > 1$ and therefore plasmon emission or absorption by the Cerenkov process is kinematically allowed in this range.

The matrix element for the helicity flipping Cerenkov process $\nu_L \rightarrow \nu_R + \gamma$ is

$$\mathcal{M} = \mu_\nu \bar{u}(p_2) \sigma^{\mu\nu} u(p_1) \sqrt{Z_l} \epsilon_\mu(k, \lambda). \quad (2)$$

¹email:mohanty@prl.ernet.in

²email:sarira@prl.ernet.in




Figure 1: Plasmon refractive index in a supernova core as a function of frequency

The photon polarisation sum in the medium for the longitudinal part is given by

$$\sum_{\lambda} \epsilon_{\mu}^L \epsilon_{\nu}^{L*} = \left(1 - \frac{1}{n^2}\right) u_{\mu} u_{\nu} + \frac{(u_{\mu} k_{\nu} + k_{\mu} u_{\nu})}{n^2 w}, \quad (3)$$

where u_{μ} is the four velocity of the medium. The rate of energy emission by this process is

$$\dot{S} = \frac{\mu_{\nu}^2}{16\pi E_1^2} \int (E_1 - w) w^2 dw Z_l f(n) (w - 2E_1)^2 \quad (4)$$

where $f(n) = (n^2 - 1)^2 / n^2$, the wave function renormalisation factor

$$Z_l = \frac{2}{n^2 + \frac{3m_{\gamma}^2}{w^2} - 1}, \quad (5)$$

and $m_{\gamma} = e^2 \tilde{\mu}^2 / 3\pi^2$ is the photon mass in the medium. The neutrino luminosity is given by

$$Q_{\nu_R} = V \int_0^{\infty} E_1^2 dE_1 f_{\nu}(E_1) \dot{S}(E_1), \quad (6)$$

where $f_{\nu}(E_1)$ is the distribution function of the neutrino.

Taking the SN core volume to be $V = 4.19 \times 10^{18} \text{ cm}^3$, the electron chemical potential $\tilde{\mu}_e = 280 \text{ MeV}$ and the neutrino chemical potential $\tilde{\mu}_{\nu} = 160 \text{ MeV}$ [5] we obtain the

neutrino luminosity to be

$$Q_{\nu_R} = 1.16 \times 10^{60} \mu_{\nu}^2 \text{ MeV}^4, \quad (7)$$

in terms of the magnetic moment of the neutrino. Assuming that the entire energy of the core collapse is not carried away by the right handed neutrinos *i.e.*, $Q_{\nu_R} < 10^{52} \text{ ergs/sec}$ we obtain the upper bound on the neutrino magnetic moment $\mu_{\nu} < 0.6 \times 10^{-11} \mu_B$. This is comparable to the bound $(0.2 - 0.8) \times 10^{-11} \mu_B$ obtained[5] from the cooling of SN1987A by the helicity flip scattering.

References

- [1] S. Mohanty and M. K. Samal, Phys. Rev. Lett. **77**, 806 (1996).
- [2] G. Raffelt, Phys. Rev. Lett. **79**, 773 (1997).
- [3] E. Bratten, Phys. Rev. Lett, **66**, 1655 (1991).
- [4] S. Sahu, Phys. Rev. **D 56**, 1688 (1997).
- [5] R. Barbieri and R. N. Mohapatra, Phys. Rev. Lett. **61**, 27 (1988).